

14P/204/5

Question Booklet No.....

(To be filled up by the candidate by blue/black ball-point pen)

Roll No.

--	--	--	--	--	--	--	--	--	--

Roll No.
(Write the digits in words)

Serial No. of OMR Answer Sheet

Day and Date (Signature of Invigilator)

INSTRUCTIONS TO CANDIDATES

(Use only blue/black ball-point pen in the space above and on both sides of the Answer Sheet)

1. Within 10 minutes of the issue of the Question Booklet, check the Question Booklet to ensure that it contains all the pages in correct sequence and that no page/question is missing. In case of faulty Question Booklet bring it to the notice of the Superintendent/Invigilators immediately to obtain a fresh Question Booklet.
2. Do not bring any loose paper, written or blank, inside the Examination Hall *except the Admit Card without its envelope.*
3. A separate Answer Sheet is given. *It should not be folded or mutilated. A second Answer Sheet shall not be provided. Only the Answer Sheet will be evaluated.*
4. Write your Roll Number and Serial Number of the Answer Sheet by pen in the space provided above.
5. **On the front page of the Answer Sheet, write by pen your Roll Number in the space provided at the top, and by darkening the circles at the bottom. Also, wherever applicable, write the Question Booklet Number and the Set Number in appropriate places.**
6. No overwriting is allowed in the entries of Roll No., Question Booklet No. and Set No. (if any) on OMR sheet and also Roll No. and OMR Sheet No. on the Question Booklet.
7. Any change in the aforesaid entries is to be verified by the invigilator, otherwise it will be taken as unfair means.
8. Each question in this Booklet is followed by four alternative answers. *For each question, you are to record the correct option on the Answer Sheet by darkening the appropriate circle in the corresponding row of the Answer Sheet, by ball-point pen as mentioned in the guidelines given on the first page of the Answer Sheet.*
9. For each question, darken only one circle on the Answer Sheet. If you darken more than one circle or darken a circle partially, the answer will be treated as incorrect.
10. *Note that the answer once filled in ink cannot be changed.* If you *do not wish to attempt* a question, leave all the circles in the corresponding row blank (such question will be awarded zero mark).
11. For rough work, use the inner back page of the title cover and the blank page at the end of this Booklet.
12. Deposit *only the OMR Answer Sheet* at the end of the Test.
13. You are not permitted to leave the Examination Hall until the end of the Test.
14. If a candidate attempts to use any form of unfair means, he/she shall be liable to such punishment as the University may determine and impose on him/her.

No. of Questions/प्रश्नों की संख्या : 150

Time/समय : 2½ Hours/घण्टे

Full Marks/पूर्णांक : 450




- Note : (1) Attempt as many questions as you can. Each question carries 3 marks. One mark will be deducted for each incorrect answer. Zero mark will be awarded for each unattempted question.

अधिकाधिक प्रश्नों को हल करने का प्रयत्न करें। प्रत्येक प्रश्न 3 अंक का है। प्रत्येक गलत उत्तर के लिए एक अंक काटा जाएगा। प्रत्येक अनुत्तरित प्रश्न का प्राप्तांक शून्य होगा।



- (2) If more than one alternative answers seem to be approximate to the correct answer, choose the closest one.

यदि एकाधिक वैकल्पिक उत्तर सही उत्तर के निकट प्रतीत हों, तो निकटतम सही उत्तर दें।

1. The graph of the derivative of f is shown in the figure below. Which of the following could be the graph of f ?



5



2. The area between the curves $y = x$ and $y = \sin x$ for $0 \leq x \leq \frac{\pi}{4}$ is
- (1) $\frac{\pi^2}{32} + \frac{1}{\sqrt{2}} - 1$ (2) $\frac{\pi^2}{32}$ (3) $\frac{\pi^2}{32} - \frac{1}{\sqrt{2}} - 1$ (4) 1
3. The center of curvature of the parabola $y^2 = 4px$ corresponding to any point on the curve is
- (1) $\left(3x - 2p, \frac{y^3}{4p^2}\right)$ (2) $\left(3x + 2p, -\frac{y^3}{4p^2}\right)$
- (3) $\left(-3x - 2p, \frac{y^3}{4p^2}\right)$ (4) $\left(3x + 2p, \frac{y^3}{4p^2}\right)$
4. The point (x, y) on the curve of $y = \sqrt{x}$ nearest to the point $(4, 0)$ is
- (1) $x = \frac{7}{2}, y = \sqrt{\frac{7}{2}}$ (2) $x = \sqrt{\frac{7}{2}}, y = \frac{7}{2}$
- (3) $x = \frac{7}{4}, y = \sqrt{\frac{7}{4}}$ (4) $x = \sqrt{\frac{6}{2}}, y = \frac{6}{2}$
5. Consider the polynomial $y = ax^n + bx^3 + c, n > 4$. The n th derivative of this polynomial $\frac{d^n y}{dx^n}$ is
- (1) $n!$ (2) n (3) $a \cdot n!$ (4) $a \cdot n$
6. If the length of a rectangle decreases at the rate of 3 cm/sec and its width increases at the rate of 2 cm/sec, the rate of change of the area of the rectangle when its length is 10 cm and its width is 4 cm is
- (1) $14 \text{ cm}^2/\text{sec}$ (2) $6 \text{ cm}^2/\text{sec}$ (3) $9 \text{ cm}^2/\text{sec}$ (4) $8 \text{ cm}^2/\text{sec}$

7. If $3x^2 + 2xy + y^2 = 2$, then the value of $\frac{dy}{dx}$ at $x = 1$ is
 (1) 2 (2) 0 (3) -2 (4) not defined
8. The function $y = x + \frac{2}{x}$ has a relative maximum at the value of x equal to
 (1) 2 (2) -2 (3) $\sqrt{2}$ (4) $-\sqrt{2}$
9. The asymptotes of the graph of the parametric equations $x = \frac{1}{t}$, $y = \frac{t}{t+1}$ are
 (1) $x = 0, y = 0$ (2) $x = 0$ only (3) $x = -1, y = 0$ (4) $x = -1$ only
10. The curvature of the cubical parabola $y = x^3$ at $(1, 1)$ is
 (1) $\frac{6}{\sqrt{1000}}$ (2) 0.3 (3) 0.6 (4) $\frac{3}{\sqrt{1000}}$
11. What is the average (mean) value of $3t^3 - t^2$ over the interval $-1 \leq t \leq 2$?
 (1) $\frac{11}{4}$ (2) $\frac{7}{2}$ (3) 8 (4) $\frac{33}{4}$
12. Consider the integral $I_n = \int x^n e^x dx$. Which of the following is true?
 (1) $I_{n+1} = x^{n+1} e^x - (n+1)I_{n-1}$ (2) $I_n = x^n e^x + nI_{n+1}$
 (3) $I_{n+1} = x^{n+1} e^x - (n+1)I_n$ (4) $I_n = x^n e^x + nI_{n-1}$
13. For what values of α and β the critical number (extremum) of the polynomial function $f(x) = x^3 + \alpha x + \beta$ is 4?
 (1) $\alpha = 48$ and $\beta = \text{arbitrary}$ (2) $\alpha = -48$ and $\beta = \text{arbitrary}$
 (3) $\alpha = \text{arbitrary}$ and $\beta = 48$ (4) $\alpha = \text{arbitrary}$ and $\beta = -48$

14. The mean value theorem guarantees the existence of a special point on the graph $y = \sqrt{x}$ between $(0, 0)$ and $(4, 2)$. What are the coordinates of this point?

(1) $(2, 1)$ (2) $(1, 1)$ (3) $(2, \sqrt{2})$ (4) None of the above

15. The value of the integral $\int_0^1 \sqrt{x^2 - 2x + 1} dx$ is

(1) -1 (2) $-\frac{1}{2}$ (3) $\frac{1}{2}$ (4) 1

16. If $U = x^y$, then

$$\frac{\left(\frac{\partial U}{\partial x} + \frac{\partial U}{\partial y}\right)}{U}$$

is

(1) $\frac{y}{x} + \log(x)$ (2) $\frac{y}{x} + \log(y)$ (3) $\frac{x}{y} + \log(x)$ (4) $\frac{x}{y} + \log(y)$

17. The slope of the line passing through the points $\left(1, -\frac{1}{2}\right)$ and $(-1, 1)$ is

(1) $\frac{3}{4}$ (2) $\frac{4}{3}$ (3) $-\frac{4}{3}$ (4) $-\frac{3}{4}$

18. The coordinates of the fourth corner of a rectangle, when three of whose corners are $(-1, 2)$, $(4, 2)$, $(-1, -3)$ is

(1) $(1, 4)$ (2) $(4, 1)$ (3) $(4, -3)$ (4) $(-1, 3)$

19. The vertex of the parabola $2y - x^2 - 4x + 6 = 0$ is

(1) $(2, 5)$ (2) $(-2, -5)$ (3) $(5, 2)$ (4) $(-5, -2)$

20. The plane P through $A(2, -3, -4)$ with normal vector $\vec{n} = 4\hat{i} - \hat{j} + 3\hat{k}$ is

(1) $4x + y + 3z = 1$ (2) $4x - 3y + z = -1$

(3) $4x - y + 3z = -1$ (4) $4x - y + 3z = 1$

11. The angle between the two planes $3x + 4y - 5z = 1$ and $4x + 5y - 6z = 1$ is
- (1) $\sin^{-1}\left(\frac{60}{\sqrt{50}}\right)$ (2) $-\cos^{-1}\left(\frac{60}{\sqrt{50}}\right)$
- (3) $\cos^{-1}\left(\frac{62}{\sqrt{50}\sqrt{77}}\right)$ (4) $-\sin^{-1}\left(\frac{60}{\sqrt{50}\sqrt{77}}\right)$
12. $M = \begin{vmatrix} 3 & 1 \\ 1 & 2 \end{vmatrix}$ is the discriminant of the conic section. Then the conic section is a
- (1) parabola (2) hyperbola
- (3) ellipse (4) rectangular hyperbola
13. The equation of the line with a slope 5 and passing through the point $(-3, 3)$ is
- (1) $y + 3 = 5(x - 3)$ (2) $y = \frac{5}{3}(x - 3)$
- (3) $y = \frac{5}{3}(x + 3)$ (4) $y - 3 = 5(x + 3)$
14. Consider two circles $x^2 + y^2 + 2ax + 2by = 0$ and $x^2 + y^2 + 2cx + 2dx = 0$ touch each other. Then the following condition is true
- (1) $ad - bc = 0$ (2) $ac - bd = 0$
- (3) $ad - bc \neq 0$ (4) No condition on a, b, c, d
15. The equation of the hyperbola with foci $(0, 0)$ and $(0, 4)$ and asymptotes $y = \pm \frac{1}{2}x$ is
- (1) $x^2 - \frac{(y-2)^2}{4} = 1$ (2) $x^2 + \frac{(y-2)^2}{4} = -1$
- (3) $y^2 - \frac{(x-2)^2}{4} = 1$ (4) $y^2 + \frac{(x-2)^2}{4} = 1$

26. Which one of the following statements is correct? The graph of $y^2 = x^2 + 9$ is symmetric about

- I The x -axis
- II The y -axis
- III The origin

- (1) I only (2) II only (3) I and II only (4) I, II and III

27. The polar form of a parabola is

- (1) $r = \frac{2p}{1 - \cos \theta}$ (2) $r = \frac{-2p}{1 - \sin \theta}$
 (3) $r = \frac{2p}{1 + \cos \theta}$ (4) $r = \frac{2p}{-1 + \sin \theta}$

28. The intervals of numbers satisfying the inequality $|x+1| > 2$ are

- (1) $x > -1$ and $x < 3$ (2) $x < 1$ and $x > -3$
 (3) $x > 3$ and $x < -1$ (4) $x > 1$ and $x < -3$

29. Solution of the inequality with absolute value $|x^2 + x - 2| < x + 3$ is

- (1) $(-\sqrt{5}, \sqrt{5})$ (2) $(-\sqrt{5}, -1) \cup (-1, \sqrt{5})$
 (3) $(-\infty, -1) \cup (-1, +\infty)$ (4) $(-1, \sqrt{5})$

30. The eigenvalues of the matrix $\sigma = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ are

- (1) $i, -i$ (2) i, i (3) $1, -1$ (4) $1, 1$

31. The inverse of the matrix

$$\begin{pmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{pmatrix}$$

is

$$(1) \frac{1}{5} \begin{pmatrix} 3 & 2 & -2 \\ 3 & 2 & 2 \\ -2 & 3 & 2 \end{pmatrix}$$

$$(2) \frac{1}{5} \begin{pmatrix} -3 & 2 & 2 \\ 2 & 3 & -2 \\ 2 & 2 & -3 \end{pmatrix}$$

$$(3) \frac{1}{5} \begin{pmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{pmatrix}$$

$$(4) \frac{1}{5} \begin{pmatrix} -3 & 2 & -2 \\ 2 & -3 & 2 \\ -2 & 2 & -3 \end{pmatrix}$$

32. The values of λ and μ for which of the following equations admit a unique solution are

$$x + y - z = 6$$

$$x + 2y + 3z = 10$$

$$x + 2y + \lambda z = \mu$$

$$(1) \lambda = 3, \mu \text{ is constant}$$

$$(2) \lambda \neq 3, \mu \text{ is arbitrary}$$

$$(3) \lambda \text{ is arbitrary, } \mu \neq 3$$

$$(4) \lambda \neq 3, \mu \text{ is rational}$$

33. The fraction $\frac{(5x+7)}{(x^2+2x-3)}$ is equal to

$$(1) \frac{2}{(x+3)} + \frac{3}{(x-1)}$$

$$(2) \frac{2}{(x-3)} + \frac{3}{(x+1)}$$

$$(3) \frac{3}{(x+3)} - \frac{2}{(x-1)}$$

$$(4) \frac{3}{(x-3)} + \frac{2}{(x+1)}$$

34. For any numbers a, b and non-zero c , if c is positive and $a < b$, then

- (1) $ac < bc$ and $\frac{a}{c} < \frac{b}{c}$ (2) $ab < ac$ and $\frac{a}{b} < \frac{a}{c}$
 (3) $ac > bc$ and $\frac{a}{c} < \frac{b}{c}$ (4) $ac < bc$ and $\frac{a}{c} > \frac{b}{c}$

35. For what values of μ the determinant of the matrix

$$A = \begin{pmatrix} 2 & -\mu & 0 \\ -1 & 5 & 1 \\ 3 & \mu^2 & 5 \end{pmatrix}$$

is 26?

- (1) (6, 2) (2) (-6, -2) (3) (-6, 2) (4) (6, -2)
36. The matrix M have three eigenvalues λ_1, λ_2 and λ_3 . One of the eigenvalues is -2 and the trace and determinant are 1 and 8 respectively. What are other two eigenvalues?
 (1) (-1, 4) (2) (1, 4) (3) (1, -4) (4) (-1, -4)
37. The function $f(x) = a_n x^n + a_{n-1} x^{n-1} + a_{n-2} x^{n-2} + \dots + a_0$ has n roots, then $(f(x))^m$, where m is a real positive integer, have
 (1) n roots (2) n^m roots (3) nm roots (4) m roots
38. If the position vector $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ and $\vec{\omega} = \omega_1\hat{i} + \omega_2\hat{j} + \omega_3\hat{k}$ is a constant vector, then $\vec{\nabla} \times \vec{\omega} \times \vec{r}$ is
 (1) ω^2 (2) $2\vec{\omega}$ (3) 0 (4) $\vec{\omega}$
39. If a force $\vec{F} = 2x^2y\hat{i} + xy\hat{j}$ displaces a particle in the xy plane from (0, 0) to (1, 4) along a curve $y = 4x^2$, then the work done is
 (1) 5 (2) 6 (3) 8 (4) 10

40. Three vectors \vec{a} , \vec{b} and \vec{c} are linearly independent if and only if

(1) $\vec{a} \cdot (\vec{b} \times \vec{c}) = 0$

(2) $\vec{a} \times (\vec{b} \times \vec{c}) = 0$

(3) $\vec{a} \cdot (\vec{b} \times \vec{c}) \neq 0$

(4) $\vec{a} \times (\vec{b} \times \vec{c}) \neq 0$

41. The moment about the point $\hat{i} + 2\hat{j} - \hat{k}$ of a force represented by $3\hat{i} + \hat{k}$ acting through the point $2\hat{i} - \hat{j} + 3\hat{k}$ is

(1) $-3\hat{i} + 11\hat{j} + 9\hat{k}$

(2) $2\hat{i} + 5\hat{j} + 2\hat{k}$

(3) $11\hat{j} + 9\hat{k}$

(4) $3\hat{i} + 11\hat{j}$

42. The value of $\vec{\nabla} \times r^n \vec{r}$, where $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ and n is an integer, is

(1) n

(2) 1

(3) 0

(4) ∞

43. $\nabla \cdot (3x^2\hat{i} + 5xy^2\hat{j} + xyz^2\hat{k})$ at the point $(1, 2, 3)$ is

(1) 36

(2) 37

(3) 38

(4) 35

44. The integral $\int_s \vec{F} \cdot \hat{n} ds$, where s is the unit sphere defined by $x^2 + y^2 + z^2 = 1$ and \vec{F} is the vector field $\vec{F} = 2x\hat{i} + y^2\hat{j} + z^2\hat{k}$, is equal to

(1) $\frac{8\pi}{3}$

(2) $\frac{\pi}{2}$

(3) $\frac{4\pi}{3}$

(4) $\frac{8\pi}{2}$

45. For the following value of m , the vectors $5\hat{i} + 6\hat{j} + 7\hat{k}$, $7\hat{i} + m\hat{j} + 9\hat{k}$ and $3\hat{i} + 20\hat{j} + 5\hat{k}$ are coplanar

(1) 8

(2) -8

(3) 6

(4) -6

46. If $\frac{dy}{dx} = e^y$ and $y = 0$ when $x = 1$, then

(1) $y = \log x$

(2) $y = \log(2 - x)$

(3) $y = -\log(2 - x)$

(4) $y = -\log x$

47. The integral $\int_C [(x^2 + xy) dx + (x^2 + y^2) dy]$, where C is the square formed by the lines $y = \pm 1, x = \pm 1$ is equal to

(1) 0

(2) 1

(3) -1

(4) ± 1

48. The general solution of $\frac{d^2y}{dx^2} + \frac{dy}{dx} - 2y = e^x$ is

(1) $y = C_1 e^x + C_2 e^{-2x} + \frac{x}{3} e^{-x}$

(2) $y = C_1 e^{-x} + C_2 e^{-2x} + \frac{x}{3} e^x$

(3) $y = C_1 e^x + C_2 e^{2x} + \frac{x}{3} e^{-x}$

(4) $y = C_1 e^{-x} + C_2 e^{2x} + \frac{x}{3} e^x$

where C_1 and C_2 are arbitrary constants.

49. The linear harmonic oscillator, $\frac{d^2x}{dt^2} + x = 0$, with the initial conditions $x(0) = 4, \dot{x}(0) = 3$ admits the solution

(1) $3 \sin t + 4 \cos t$

(2) $4 \sin t + 3 \cos t$

(3) $3 \sin t - 4 \cos t$

(4) $4 \sin t - 3 \cos t$

50. If $\frac{dy}{dx} = y \tan x$, then y is equal to

(1) $\frac{1}{2} \tan^2 x + c$

(2) $c \tan x + c$

(3) $c \sec x$

(4) $\ln |\cos x| + c$

where c is a constant.

51. If $f'(x) = -f(x)$ and $f(1) = 1$, then $f(x) = ?$

- (1) $\frac{1}{2} e^{(-2x+2)}$ (2) $e^{-(x+1)}$ (3) e^{1-x} (4) e^{-x}

52. The inverse Laplace transform of the function $\log\left(1 + \frac{w^2}{s^2}\right)$ is

- (1) $\frac{2}{t}(1 - \cos wt)$ (2) $\frac{2}{(1 - \cos wt)}$

- (3) $\frac{2}{(1 - \sin wt)}$ (4) $1 - \sin wt$

53. $C = x \frac{dy}{dx} - y^3 + x$ is the invariant (constant) curve for

- (1) $x \frac{d^2y}{dx^2} = 3y^2 \frac{dy}{dx} - 1 - \frac{dy}{dx}$ (2) $x \frac{d^2y}{dx^2} = 1 - 3y^2 \frac{dy}{dx} + \frac{dy}{dx}$

- (3) $\frac{d^2y}{dx^2} = 3y^2 \frac{dy}{dx} - 1 - \frac{dy}{dx}$ (4) $x \frac{d^2y}{dx^2} = 3y^2 \frac{dy}{dx} - 1 - x \frac{dy}{dx}$

54. The curve $y = e^x + e^{-x}$ satisfies the differential equation

- (1) $\frac{dy}{dx} = y$ (2) $\frac{dy}{dx} = -y$ (3) $\frac{d^2y}{dx^2} = y$ (4) $\frac{d^2y}{dx^2} = -y$

55. For the differential equation $y \frac{dy}{dx} + 2 \cos(y) y = 1$, which of the following is true?

- (1) The differential equation is first-order linear and homogenous
 (2) The differential equation is first-order linear and non-homogenous
 (3) The differential equation is first-order nonlinear and homogenous
 (4) The differential equation is first-order nonlinear and non-homogenous

56. If $\tan a = \frac{1}{3}$ and $\tan b = \frac{1}{2}$, then $a + b$ is
- (1) $\frac{3\pi}{2}$ (2) $\frac{3\pi}{4}$ (3) $\frac{\pi}{2}$ (4) $\frac{\pi}{4}$
57. $\arcsin r = \theta$, then θ is
- (1) $-i \ln(\sqrt{1-r^2} + ir)$ (2) $i \ln(\sqrt{1-r^2} + ir)$
 (3) $-i \ln(\sqrt{1-r^2} - ir)$ (4) $i \ln(\sqrt{1-r^2} - ir)$
58. Which of the following defines a function f for which $f(-x) = -f(x)$?
- (1) $f(x) = x^2$ (2) $f(x) = \sin x$
 (3) $f(x) = \cos x$ (4) $f(x) = e^x$
59. If $\log(a+ib) = (c+id)$, then
- (1) $c = \log(\sqrt{a^2+b^2})$, $d = \tan^{-1}\left(\frac{b}{a}\right)$ (2) $c = \log(a^2+b^2)$, $d = \tan^{-1}\left(\frac{a}{b}\right)$
 (3) $c = \log(\sqrt{a^2+b^2})$, $d = \frac{1}{2} \tan^{-1}\left(\frac{b}{a}\right)$ (4) $c = \log(a^2+b^2)$, $d = \tan^{-1}\left(\frac{b}{a}\right)$
60. The real value of the function $\frac{F(z_1)}{F(z_2)}$ for $F(z) = z + |z|^2$, $z_1 = 3e^{i\frac{\pi}{2}}$ and $z_2 = 2e^{\pi}$ is
- (1) 3 (2) 0 (3) -2 (4) 1
61. Let the functions f and g have 6 and 3 roots, respectively. If all the roots of g are also roots of f , then how many roots does the function $f \times g$ has?
- (1) 3 (2) 6 (3) 9 (4) 18

62. One of the values of $(i)^{\frac{1}{3}}$ is
 (1) $-i$ (2) i (3) -1 (4) 1
63. For a right angled triangle if one of the angle is α ($\alpha \neq \frac{\pi}{2}$), the other angle is
 (1) $\pi - \alpha$ (2) $\alpha - \frac{\pi}{2}$ (3) $\frac{\pi}{2} - \alpha$ (4) $\frac{\pi}{2} - \alpha$
64. Consider the matrix $A(\theta) = \begin{pmatrix} \sin(\theta) & -\cos(\theta) \\ \cos(\theta) & \sin(\theta) \end{pmatrix}$ and $\rho = |A(\theta)| |A(\phi)|$. For $\phi = \theta - \frac{\pi}{2}$ the value of ρ is
 (1) 2 (2) 1 (3) 0 (4) -1
65. The points of intersection of $f = \sin^2(2\theta)$ and $f = \cos^2(2\theta)$ between $\frac{-\pi}{2}$ to $\frac{\pi}{2}$ are
 (1) $\left(\frac{-\pi}{4}, \frac{\pi}{4}\right)$ (2) $\left(\frac{-3\pi}{8}, \frac{3\pi}{8}\right)$ (3) $\left(\frac{-\pi}{8}, \frac{\pi}{8}\right)$ (4) $\left(\frac{-3\pi}{8}, \frac{3\pi}{8}\right)$
66. A gun moving at a speed of 30 m/sec fires at an angle 30° with a velocity 150 m/s relative to the gun. The distance between the gun and the projectile when projectile hits the ground ($g = 10$ m/sec) is
 (1) 1850 m (2) 1750 m (3) 1950 m (4) 1050 m
67. The displacement of particle executing simple harmonic motion obeys the equation $y = 1.60 \sin(1.3t)$. Here, y is in centimetres and t is in seconds. The magnitude of the velocity at $t = 0$ is
 (1) $v = 1.08$ m/s (2) $v = 0.08$ m/s (3) $v = 3.08$ m/s (4) $v = 2.08$ m/s

68. One spring has force constant 200 Nm^{-1} , another has force constant 500 Nm^{-1} . If they are joined in series, the force constant will be nearest to
- (1) 700 N/m (2) 300 N/m (3) 143 N/m (4) 100 N/m
69. A particle moves in a straight line so that its distance at time t from a fixed point of the line is $8t - 3t^2$. What is the total distance covered by the particle between $t = 1$ and $t = 2$?
- (1) 1 (2) $\frac{4}{3}$ (3) $\frac{5}{3}$ (4) 2
70. The degrees of freedom of the particle constrained to move only on surface of the sphere is
- (1) 2 (2) 3 (3) 0 (4) 1
71. A body whose three principal moments of inertia are all equal, that is $I_1 = I_2 = I_3$, is called as
- (1) asymmetrical top (2) symmetrical top
(3) spherical top (4) None of the above
72. What is the necessary condition for a force \vec{F} to be conservative?
- (1) $\vec{\nabla} \cdot \vec{F} = 0$ (2) $\vec{\nabla} \times \vec{F} = 0$ (3) $\vec{\nabla} \cdot \vec{F} \neq 0$ (4) $\vec{\nabla} \times \vec{F} \neq 0$
73. When a rigid body rotates about an axis and the external torque is zero, then for that body the following is a constant
- (1) Angular velocity (2) Moment of inertia
(3) Linear momentum (4) Angular momentum

74. If a body has mass m , velocity at centre of mass v_c , moment of inertia I_c and rotational velocity ω , then total kinetic energy is

- (1) $\frac{1}{2}mv_c^2$ (2) $\frac{1}{2}I_c\omega^2$
 (3) $\frac{1}{2}I_c\omega^2 - \frac{1}{2}mv_c^2$ (4) $\frac{1}{2}I_c\omega^2 + \frac{1}{2}mv_c^2$

75. The angular momentum of a rotational body, with angular velocity ω and moment of inertia I , is given by

- (1) $\frac{1}{2}I\omega$ (2) $I\omega$ (3) $\frac{1}{2}I\omega^2$ (4) $I\omega^2$

76. In metals the skin depth for electromagnetic waves

- (1) increases with increase in frequency
 (2) decreases with increase in frequency
 (3) does not depend on frequency
 (4) increases or decreases with frequency depending on the conductivity of metal

77. A plane polarized electromagnetic wave with \vec{E} vector parallel to the plane of incidence is incident from air to glass. It is found that $\theta_i + \theta_t = 90^\circ$, where θ_i is the angle of incidence and θ_t is the angle of transmittance then

- (1) there will not be any reflected wave
 (2) the reflected wave will be in a direction perpendicular to transmitted wave
 (3) the reflected wave will be in a direction perpendicular to incident wave
 (4) the reflected wave will be perpendicular to the refracted wave

78. The average value of the Poynting vector for a plane polarized sinusoidal electromagnetic wave in free space is given by

- (1) $\frac{1}{2}\epsilon_0 E^2$ (2) $\frac{1}{2}\mu_0 B_0^2$ (3) $\frac{1}{2}\frac{\mu_0 B^2}{C}$ (4) $\frac{1}{2}C\epsilon_0 E_0^2$

E_0 and B_0 are the peak values of the amplitudes of electric and magnetic field.

79. The dielectric constant of any dielectric materials for electromagnetic waves

- (1) increases with frequency
 (2) is independent of frequency
 (3) decreases with frequency
 (4) decreases with frequency in radio frequency range but increases with frequency in optical range

80. How many 2 input NAND gates will be required to realize the operation of 3 input OR gates?

- (1) 3 (2) 4 (3) 5 (4) 6

81. The simplified Boolean expression in POS for the Boolean expression $Y = ABC\bar{C} + A\bar{B}C + \bar{A}BC + ABC$ is given by

- (1) $Y = AB + BC + AC$ (2) $Y = (A+B) \cdot (B+C) \cdot (C+A)$
 (3) $Y = (\bar{A} + B) \cdot (\bar{B} + C) \cdot (\bar{C} + A)$ (4) $Y = \bar{A}B + \bar{B}C + \bar{C}A$

82. Two 4-bit numbers can be added by using

- (1) 4 full adders (2) 8 half adders
 (3) 3 full adder and 1 half adder (4) 1 full adder and 3 half adder

83. Which one of the following is not the basic logic gate?
- (1) AND (2) OR (3) NOT (4) XOR
84. The wavelength of an electromagnetic wave of frequency 10 GHz travelling in a medium with $\mu = 4\pi \times 10^{-7} \text{ H/m}$ and $\epsilon = \frac{1}{36\pi} \times 10^{-9} \text{ F/m}$ will be
- (1) 3 cm (2) 3 metre (3) 30 cm (4) 30 metre
85. A material has $\sigma = 10^{-2} \text{ s/m}$ and $\epsilon = 2\epsilon_0$ at what frequency will the conduction current be equal to the displacement current?
- (1) $6.3 \times 10^6 \text{ Hz}$ (2) $9.1 \times 10^7 \text{ Hz}$
 (3) $3.1 \times 10^8 \text{ Hz}$ (4) $5.3 \times 10^9 \text{ Hz}$
86. Which one of the following is not a Maxwell's equation of electromagnetic?
- (1) $\oint_s \vec{D} \cdot d\vec{s} = q$ (2) $\oint \vec{B} \cdot d\vec{l} = \mu_0 I$
 (3) $\oint \vec{H} \cdot d\vec{l} = \int_s \left(\vec{J} + \frac{\partial \vec{D}}{\partial t} \right) \cdot d\vec{s}$ (4) $\oint \vec{E} \cdot d\vec{l} = -\frac{\partial}{\partial t} \int_s \vec{B} \cdot d\vec{s}$
87. For plane electromagnetic waves in vacuum which of the following statements is not true?
- (1) These are transverse in nature
 (2) Electric and magnetic field waves are in phase
 (3) There is a phase difference of 90° between electric and magnetic fields
 (4) $\vec{E} \times \vec{H}$ points in the direction of propagation of electromagnetic wave

88. The de Broglie wavelength of an electron moving with velocity 10 m/sec is (given $h = 6.63 \times 10^{-34}$ J-sec, $m_e = 9.1 \times 10^{-31}$ kg)
- (1) 3.6×10^{-11} m (2) 1.44×10^{-10} m
 (3) 11.0×10^{-11} m (4) 7.3×10^{-11} m
89. If we pour some drops of water between the plate and lens in Newton's ring experiment, then the rings will
- (1) increase in diameter (2) decrease in diameter
 (3) become elliptical (4) become invisible
90. Two polarizing sheets have directions such that the transmitted light has maximum intensity I_{\max} . Through what angle must either sheet be turned so that the intensity of transmitted light becomes $\frac{I_{\max}}{2}$?
- (1) $\pm 30^\circ$ (2) $\pm 60^\circ$ (3) $\pm 45^\circ$ (4) $\pm 90^\circ$
91. A circularly polarised light can be distinguished from unpolarized light by passing it through
- (1) Nicol prism (2) polarizing sheet
 (3) half-wave plate (4) quarter-wave plate
92. If mirror M_2 in Michelson interferometer is moved through 0.233 mm, then 792 fringes are counted. The wavelength of light is
- (1) 715 nm (2) 656 nm (3) 588 nm (4) 536 nm

98. If C_p and C_v are the molar specific heats of a gas at constant pressure and constant volume respectively. The ratio of adiabatic and isothermal moduli of elasticity will be
- (1) $\frac{C_p - C_v}{C_p}$ (2) $\frac{C_p - C_v}{C_v}$ (3) $\frac{C_v}{C_p}$ (4) $\frac{C_p}{C_v}$
99. A Carnot engine has an efficiency of 40% and a heat sink temperature of 27 °C. What should be the temperature of heat sink so that the efficiency becomes 50%?
- (1) 200 °K (2) 250 °K (3) 325 °K (4) 350 °K
100. Total time of flight of a projectile launched with velocity u at angle θ with the horizontal is
- (1) $\frac{u \sin \theta}{g}$ (2) $\frac{2u \cos \theta}{g}$ (3) $\frac{2u \sin \theta}{g}$ (4) $\frac{u \cos \theta}{g}$
101. A bullet of mass m travelling with velocity v gets embedded into a sand bag of mass M suspended by an instretchable string. The loss of kinetic energy in the process would be
- (1) $\frac{1}{2} \frac{m^2 v^2}{(M+m)}$ (2) $\frac{1}{2} \frac{M^2 v^2}{(M+m)}$ (3) $\frac{1}{2} \frac{(M+m)^2 v^2}{m}$ (4) $\frac{1}{2} \frac{mM}{(m+M)} v^2$
102. The largest and the smallest distance of the earth from the sun in its orbit are r_1 and r_2 respectively. Its distance from the sun at the perpendicular to the major axis of the orbit passing through the sun would be
- (1) $\frac{2r_1 r_2}{(r_1 + r_2)}$ (2) $\frac{(r_1 + r_2)}{2r_1 r_2}$ (3) $\frac{r_1 + r_2}{2}$ (4) $\frac{r_1 - r_2}{2}$
103. If the noise level in Varanasi is 80 dB and that in Chandigarh is 40 dB, then the intensity of noise in Varanasi exceeds that in Chandigarh by a factor of
- (1) 2 (2) 2⁴ (3) 10⁴ (4) 20

104. When an intense beam of laser light goes from air into water there is no change in its
 (1) intensity (2) frequency (3) velocity (4) wavelength
105. The magnification of the image by a concave mirror of focal length f is m . If the image is real the distance of the object from the mirror would be
 (1) $(m-1)f$ (2) $(m+1)f$ (3) $\frac{m+1}{m}f$ (4) $\left(\frac{m-1}{m}\right)f$
106. If the half-life of a radio active substance is 3 days, then by what factor would its activity reduce in 9 days?
 (1) $\frac{1}{3}$ (2) $\frac{2}{3}$ (3) $\frac{1}{8}$ (4) $\frac{7}{8}$
107. A system of three identical condensers will store maximum energy of
 (1) two are connected in series and third in parallel to them
 (2) two are connected in parallel and third in series with the combination
 (3) all three connected in series
 (4) all three connected in parallel
108. A mass spectrograph is used for the determination of
 (1) specific charge of an ion (2) atomic mass
 (3) spectral lines of isotopes (4) atomic charge
109. The series of spectral lines in the spectrum of hydrogen atom that lies partly in the ultraviolet and partly in the visible region is called
 (1) Balmer series (2) Lyman series
 (3) Brackett series (4) Paschen series

110. Neglecting the relativistic effect the wavelength associated with electron of kinetic energy E is proportional to

- (1) \sqrt{E} (2) $\frac{1}{\sqrt{E}}$ (3) E^2 (4) $\frac{1}{E^2}$

111. The unit for measurement of man's exposure to nuclear radiation is

- (1) Curie (2) Becquerel (3) Rutherford (4) Fermi

112. A metallic wire of length L hanging from the roof is stretched by an small amount l when a body of mass m is attached to its free end. The mechanical energy stored in the wire is

- (1) $\frac{mgl}{L}$ (2) $\frac{mgl^2}{L}$ (3) $\frac{mgl}{2}$ (4) $\frac{mgl^2}{2L}$

113. Two uniform circular discs A and B of equal masses and thicknesses are made of materials of densities ρ_A and ρ_B respectively. If their moment of inertia about an axis passing through the center and normal to the circular faces are I_A and I_B respectively, then

- (1) $\frac{I_A}{I_B} = \frac{\rho_A}{\rho_B}$ (2) $\frac{I_A}{I_B} = \frac{\rho_B}{\rho_A}$ (3) $\frac{I_A}{I_B} = \left(\frac{\rho_A}{\rho_B}\right)^2$ (4) $\frac{I_A}{I_B} = \left(\frac{\rho_B}{\rho_A}\right)^2$

114. The main use of a voltage series negative feedback amplifier is as a

- (1) power amplifier
 (2) current amplifier
 (3) impedance matching device
 (4) low input impedance voltage amplifier

115. In RC coupled transistor amplifier the upper cut off in frequency response is obtained due to
- (1) coupling capacitance (2) blocking capacitance
(3) by pass capacitance (4) junction capacitance
116. The width of the depletion region layer of a P-N junction diode
- (1) decreases with increasing doping concentration
(2) increases with increasing doping concentration
(3) is independent of doping concentration
(4) decrease with increasing reverse bias
117. Avalanche break down in Zener diode is a phenomena primarily caused by ionization of immobile ions
- (1) due to high electric field
(2) due to collision with high velocity minority charge carriers
(3) due to collision with high velocity majority charge carriers
(4) due to tunnelling of charge carriers
118. A circuit having an inductance of $\frac{1}{\pi}$ Henry and resistance of 100 ohms is connected to AC power supply at 50 Hz frequency. The reactance and impedance of the circuit is
- (1) 100 Ω , 100 Ω (2) 141.1 Ω , 100 Ω
(3) 100 Ω , 141.1 Ω (4) 141.1 Ω , 141.1 Ω

130. X-rays of 10.0 p.m. are scattered from a target in all directions and the maximum wavelength present in the scattered X-rays is 14.9 p.m. Find the wavelength of the X-rays scattered at 45°

- (1) 10.7 p.m. (2) 12.425 p.m. (3) 11.25 p.m. (4) 9.3 p.m.

131. The wave particle duality was demonstrated by the

- (1) Stern-Gerlach experiment (2) Davisson-Germer experiment
(3) Franck-Hertz experiment (4) Michelson-Morley experiment

132. Raman scattering is a quantum mechanical process involving

- (1) one photon (2) one photon and one electron
(3) two photons (4) two photon and one electron

133. Planck's radiation formula reduces to

- (1) Rayleigh-Jeans formula at low frequencies
(2) Rayleigh-Jeans formula high frequencies
(3) Wien's displacement formula at low temperature
(4) Rayleigh-Jeans formula at low temperature

134. Two linearly polarised light waves of unequal amplitudes with their planes of polarization perpendicular to each other on superposition give rise to

- (1) circularly polarized light (2) plane polarised light
(3) unpolarised light (4) elliptically polarized light

140. For a van der Waals' gas the Joule-Thomson coefficient is given by

- (1) $\frac{1}{C_p} \left[b - \frac{2a}{RT} \right]$ (2) $\frac{1}{C_v} \left[\frac{2a}{RT} - b \right]$
 (3) $\frac{1}{C_p} \left[\frac{2a}{RT} - b \right]$ (4) $\frac{1}{C_v} \left[b - \frac{2a}{RT} \right]$

141. The Fourier series

$$F(x) = \frac{3}{2} + \frac{6}{\pi} \left[\sin \frac{\pi x}{5} + \frac{1}{3} \sin \frac{3\pi x}{5} + \frac{1}{5} \sin \frac{5\pi x}{5} + \dots \right]$$

represents a square wave of

- (1) amplitude 3 and time period 5 (2) amplitude $\frac{3}{2}$ and time period 10
 (3) amplitude 3 and time period 10 (4) amplitude $\frac{3}{2}$ and time period 5
142. A reversible heat engine converts $\frac{1}{6}$ th heat, which it absorbs from source into useful work. When the temperature of the sink is reduced by 60°C , its efficiency is doubled. Then the temperature of the source is
- (1) 240 K (2) 300 K (3) 480 K (4) 360 K

143. A diffraction grating is illuminated by a Laser light of wavelength 500 nm. If the second order spectral line is observed at 30° , then the number of lines per centimetre of grating is

- (1) 5000 (2) 6000 (3) 4000 (4) 3000
144. For a series L-C-R resonance circuit the power factor at resonance is
- (1) infinity (2) zero (3) half (4) unity

145. A bridge rectifier is preferred over an ordinary full-wave rectifier because
- (1) its rectification efficiency is high
 - (2) its ripple factor is small
 - (3) its transformer does not require center tap secondary
 - (4) its peak inverse voltage is low
146. Indicate the false statement regarding the early effect in transistor.
- (1) base current decreases with increasing $|V_{CB}|$
 - (2) emitter current increase with increase $|V_{CB}|$
 - (3) α decreases with increasing $|V_{CB}|$
 - (4) β increases with increasing $|V_{CB}|$
147. What will be the maximum wave length of light that will cause the photoelectrons to be emitted from sodium target whose work function is 23 eV ($h = 4.14 \times 10^{-15}$ eV \times sec) ?
- (1) 270 nm (2) 675 nm (3) 810 nm (4) 540 nm
148. Indicate the false statement about the conclusions drawn from Michelson-Morley experiment
- (1) hypothetical ether does not exist
 - (2) all motions are relative to a universal frame of reference
 - (3) the speed of light is same for all observers
 - (4) all motions are relative to a specified frame of reference

149. Gibbs' free energy G is defined as

(1) $G = u + PV + TS$

(2) $u - PV + TS$

(3) $G = u + PV - TS$

(4) $u - PV - TS$

150. The radius of gyration of a thin uniform rod of mass $M = 100$ gm and length $l = 1$ metre about an axis passing through its center of gravity and perpendicular of its length is

(1) $k = \frac{1}{2\sqrt{3}}$ metre

(2) $k = \frac{1}{3\sqrt{3}}$ metre

(3) $k = \frac{1}{4\sqrt{3}}$ metre

(4) $k = \frac{1}{6\sqrt{3}}$ metre

अभ्यर्थियों के लिए निर्देश

(इस पुस्तिका के प्रथम आवरण-पृष्ठ पर तथा उत्तर-पत्र के दोनों पृष्ठों पर केवल नीली या काली बाल-प्वाइंट पेन से ही लिखें।)

1. प्रश्न पुस्तिका मिलने के 10 मिनट के अन्दर ही देख लें कि प्रश्नपत्र में सभी पृष्ठ मौजूद हैं और कोई प्रश्न छूटा नहीं है। पुस्तिका दोषयुक्त पाये जाने पर इसकी सूचना तत्काल कक्ष-निरीक्षक को देकर सम्पूर्ण प्रश्नपत्र की दूसरी पुस्तिका प्राप्त कर लें।
2. परीक्षा भवन में लिफाफा रहित प्रवेश-पत्र के अतिरिक्त, लिखा या सादा कोई भी खुला कागज साथ में न लायें।
3. उत्तर-पत्र अलग से दिया गया है। इसे न तो मोड़ें और न ही विकृत करें। दूसरा उत्तर-पत्र नहीं दिया जायेगा, केवल उत्तर-पत्र का ही मूल्यांकन किया जायेगा।
4. अपना अनुक्रमांक तथा उत्तर-पत्र का क्रमांक प्रथम आवरण-पृष्ठ पर पेन से निर्धारित स्थान पर लिखें।
5. उत्तर-पत्र के प्रथम पृष्ठ पर पेन से अपना अनुक्रमांक निर्धारित स्थान पर लिखें तथा नीचे दिये वृत्तों को गाढ़ा कर दें। जहाँ-जहाँ आवश्यक हो वहाँ प्रश्न-पुस्तिका का क्रमांक तथा सेट का नम्बर उचित स्थानों पर लिखें।
6. ओ० एम० आर० पत्र पर अनुक्रमांक संख्या, प्रश्न-पुस्तिका संख्या व सेट संख्या (यदि कोई हो) तथा प्रश्न-पुस्तिका पर अनुक्रमांक सं० और ओ० एम० आर० पत्र सं० की प्रविष्टियों में उपरिलेखन की अनुमति नहीं है।
7. उपर्युक्त प्रविष्टियों में कोई भी परिवर्तन कक्ष निरीक्षक द्वारा प्रमाणित होना चाहिये अन्यथा यह एक अनुचित साधन का प्रयोग माना जायेगा।
8. प्रश्न-पुस्तिका में प्रत्येक प्रश्न के चार वैकल्पिक उत्तर दिये गये हैं। प्रत्येक प्रश्न के वैकल्पिक उत्तर के लिये आपको उत्तर-पत्र की सम्बन्धित पंक्ति के सामने दिये गये वृत्त को उत्तर-पत्र के प्रथम पृष्ठ पर दिये गये निर्देशों के अनुसार पेन से गाढ़ा करना है।
9. प्रत्येक प्रश्न के उत्तर के लिये केवल एक ही वृत्त को गाढ़ा करें। एक से अधिक वृत्तों को गाढ़ा करने पर अथवा एक वृत्त को अपूर्ण भरने पर वह उत्तर गलत माना जायेगा।
10. ध्यान दें कि एक बार स्याही द्वारा अंकित उत्तर बदला नहीं जा सकता है। यदि आप किसी प्रश्न का उत्तर नहीं देना चाहते हैं, तो सम्बन्धित पंक्ति के सामने दिये गये सभी वृत्तों को खाली छोड़ दें। ऐसे प्रश्नों पर शून्य अंक दिये जायेंगे।
11. रफ़ कार्य के लिये प्रश्न-पुस्तिका के मुखपृष्ठ के अन्दर वाले पृष्ठ तथा अंतिम पृष्ठ का प्रयोग करें।
12. परीक्षा के उपरान्त केवल ओ०एम०आर० उत्तर-पत्र परीक्षा भवन में जमा कर दें।
13. परीक्षा समाप्त होने से पहले परीक्षा भवन से बाहर जाने की अनुमति नहीं होगी।
14. यदि कोई अभ्यर्थी परीक्षा में अनुचित साधनों का प्रयोग करता है, तो वह विश्वविद्यालय द्वारा निर्धारित दंड का/की, भागी होगा/होगी।